

## East Africa leDEA

### Funding acknowledgement-Adult and Pediatric

Research reported in this publication was supported by the National Institute Of Allergy And Infectious Diseases (NIAID), Eunice Kennedy Shriver National Institute Of Child Health & Human Development (NICHD), National Institute On Drug Abuse (NIDA), National Cancer Institute (NCI), and the National Institute of Mental Health (NIMH), in accordance with the regulatory requirements of the National Institutes of Health under Award Number U01AI069911East Africa leDEA Consortium. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

### Site investigators and cohorts (without data managers):

Lameck Diero, Samuel Ayaya, MOI University, AMPATH Plus, Eldoret, Kenya; Elizabeth Bukusi, KEMRI (Kenya Medical Research Institute), Kisumu, Kenya; John Ssali, Masaka Regional Referral Hospital, Masaka, Uganda; Mwebesa Bosco Bwana, Mbarara University of Science and Technology (MUST), Mbarara, Uganda; Barbara Castelnuovo, Infectious Diseases Institute (IDI), Mulago, Uganda; Fred Nalugoda, Rakai Health Sciences Program, Kalisizo, Uganda; G.R. Somi, NACP (National AIDS Control Program) Dar es Salaam, Tanzania; Rita Elias Lyamuya, Morogoro Regional Hospital, Morogoro, Tanzania; Kapella Ngonyani, Tumbi Regional Hospital, Pwani, Tanzania; Mark Urassa, Denna Michael, National Institute for Medical Research (NIMR), Kisesa HDSS, Mwanza, Tanzania; Kara Wools-Kaloustian, Constantin Yiannoutsos, Rachel Vreeman, Indiana University School of Medicine, Indiana University, Indianapolis, IN, USA; Batya Elul, Columbia University, New York City, NY, USA; Jennifer Syvertsen, Ohio State University, Columbus, OH, USA; Rami Kantor, Brown University/Miriam Hospital, Providence, RI, USA; Jeffrey Martin, Craig Cohen, University of California, San Francisco, CA, USA; Paula Braitstein, University of Toronto, Toronto, Canada

### Site investigators and cohorts (with data managers):

Diero L, Ayaya S, Sang E, MOI University, AMPATH Plus, Eldoret, Kenya; Bukusi E, Charles Karue Kibaara, Elisheba Mutegi, KEMRI (Kenya Medical Research Institute), Kisumu, Kenya; John Ssali, Mathew Ssemakadde, Masaka Regional Referral Hospital, Masaka, Uganda; Mwebesa Bosco Bwana, Michael Kanyesigye, Mbarara University of Science and Technology (MUST), Mbarara, Uganda; Barbara Castelnuovo; John Michael Matovu, Infectious Diseases Institute (IDI), Mulago, Uganda; Fred Nalugoda, Francis X. Wasswa, Rakai Health Sciences Program, Kalisizo, Uganda; G.R. Somi, Joseph Nondi, NACP (National AIDS Control Program) Dar es Salaam, Tanzania; Rita Elias Lyamuya, Francis Mayanga, Morogoro Regional Hospital, Morogoro, Tanzania; Kapella Ngonyani, Jerome Lwali, Tumbi Regional Hospital, Pwani, Tanzania; Mark Urassa, Denna Michael, Richard Machemba, National Institute for Medical Research (NIMR), Kisesa HDSS, Mwanza, Tanzania; Kara Wools-Kaloustian, Constantin Yiannoutsos, Rachel Vreeman, Beverly Musick, Indiana University School of Medicine, Indiana University, Indianapolis, IN, USA; Batya Elul, Columbia University, New York City, NY, USA; Jennifer Syvertsen, Ohio State University, Columbus, OH, USA; Rami Kantor, Brown University/Miriam Hospital, Providence, RI, USA; Jeffrey Martin, Megan Wenger, Craig Cohen, Jayne Kulzer, University of California, San Francisco, CA, USA; Paula Braitstein, University of Toronto, Toronto, Canada